

# Vav family exchange factors: an integrated regulatory and functional view

Xosé R Bustelo\*

Centro de Investigación del Cáncer and Instituto de Biología Molecular y Celular del Cáncer; Consejo Superior de Investigaciones Científicas (CSIC) and University of Salamanca; Campus Unamuno; Salamanca, Spain

**Keywords:** cancer, cardiovascular biology, Dbl-homology, disease, GDP/GTP exchange factors, immunology, nervous system, Rho GTPases, signaling, therapies, Vav

**Abbreviations:** Ac, acidic; Ahr, aryl hydrocarbon receptor; cGMP, cyclic guanosine monophosphate; CH, calponin homology; CSH3, most C-terminal SH3 domain of Vav proteins; DAG, diacylglycerol; DH, Dbl-homology domain; GEF, guanosine nucleotide exchange factor; HIV, human immunodeficiency virus; IP<sub>3</sub>, inositoltriphosphate; NFAT, nuclear factor of activated T-cells; NSH3, most N-terminal SH3 domain of Vav proteins; PH, plekstrin-homology domain; Phox, phagocyte oxidase; PI3K, phosphatidylinositol-3 kinase; PIP<sub>3</sub>, phosphatidylinositol (3,4,5)-triphosphate; PKC, protein kinase C; PKD, protein kinase D; PLC-g, phospholipase C-g; PRR, proline-rich region; PTK, protein tyrosine kinase; SH2, Src homology 2; SH3, Src homology 3; SNP, single nucleotide polymorphism; TCR, T-cell receptor; ZF, zinc finger region.

The Vav family is a group of tyrosine phosphorylation-regulated signal transduction molecules hierarchically located downstream of protein tyrosine kinases. The main function of these proteins is to work as guanosine nucleotide exchange factors (GEFs) for members of the Rho GTPase family. In addition, they can exhibit a variety of catalysis-independent roles in specific signaling contexts. Vav proteins play essential signaling roles for both the development and/or effector functions of a large variety of cell lineages, including those belonging to the immune, nervous, and cardiovascular systems. They also contribute to pathological states such as cancer, immune-related dysfunctions, and atherosclerosis. Here, I will provide an integrated view about the evolution, regulation, and effector properties of these signaling molecules. In addition, I will discuss the pros and cons for their potential consideration as therapeutic targets.

## Introduction

The first member of the Vav family was discovered due to an artifactual oncogenic activation of the *VAV1* gene during the course of focus formation assays conducted in M. Barbacid's lab.<sup>1</sup> Since it was the sixth human oncogene identified by that group, it was designated as the sixth letter of the Hebrew alphabet (VAV). Mouse and human *Vav2* and *Vav3* genes were isolated using standard cloning procedures in the 1995–1999 period.<sup>2–4</sup> The rest of family members were identified during the extensive


characterization of genomes from multiple species that took place during this last decade. The isolation of the other family members led to the progressive designation of the founding member of the family as Vav1, a name that has become common currency in the field. Despite such a serendipitous discovery, we now know that Vav proteins play crucial signaling roles in a large variety of organisms and cell types. They also show a quite idiosyncratic activation by direct tyrosine phosphorylation, a property that makes them critical elements in protein tyrosine kinase (PTK)-regulated pathways. Consistent with the oncogenic activity shown by the founder member of the family, recent data have revealed that these proteins also play roles in tumorigenesis, metastasis, and many other diseases. In this review, I will provide a general overview about this family, including new data about its evolution, regulation, function, signaling specificity, and potential therapeutic value. Readers can find additional information about Vav proteins in previous review articles,<sup>5–10</sup> book chapters,<sup>11</sup> and online resources.<sup>12–14</sup>

## Phylogenetic Distribution and Structure

The earliest evidence of the presence of a Vav family protein in the phylogenetic tree is found in Choanoflagellates, a group of unicellular aquatic protists considered to be the closest relatives to metazoans.<sup>15</sup> They are also the first known organisms that developed PTK- and Src homology 2 (SH2)-encoding genes,<sup>15</sup> thus emphasizing the close link that exists between Vav proteins and PTK-driven signal transduction events. Since then, members of this family have been found in all animal metazoans characterized so far. Single Vav family members are present in invertebrate species while vertebrates contain 3 family members. However, alternative splicing events yield different protein isoforms in all species. In mammals, Vav1 is primarily detected in haematopoietic cells whereas Vav2 and Vav3 display broader expression patterns.<sup>2–4,16,17</sup>

\*Correspondence to: Xosé R Bustelo; Email: [xbustelo@usal.es](mailto:xbustelo@usal.es)  
Submitted: 07/27/2014; Revised: 09/02/2014; Accepted: 09/05/2014  
<http://dx.doi.org/10.4161/21541248.2014.973757>

This is an Open Access article distributed under the terms of the Creative Commons Attribution-Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited. The moral rights of the named author(s) have been asserted.



**Figure 1.** Examples of the multidomain structure of some Vav family proteins. Abbreviations for domains have been indicated in the main text. The shape of the DH tries to mimic the 3 dimensional structure of this domain. Phosphorylation sites are shown as yellow circles. Amino acid numbers correspond to the primary sequence of mouse Vav1. An Ac  $\alpha$  helix involved in the stabilization of the autoinhibited structure of Vav proteins is shown as a blue box. Cel, *C. elegans*; Dm, *D. melanogaster*.

phosphotyrosine binding SH2 domain. As will be discussed later, these structural idiosyncrasies have a significant impact in the regulation and function of these proteins.

## Signaling Roles

Tyrosine-phosphorylated Vav proteins exhibit both catalysis-dependent and independent activities during cell signaling (Fig. 2). The enzyme activity involves the stimulation of GDP/GTP exchange on

All Vav family proteins contain a basic, evolutionarily conserved structural scaffold composed of an N-terminal calponin-homology (CH) domain, an acidic (Ac) region, a catalytic Dbl-homology (DH) domain involved in the guanosine nucleotide exchange reaction, a pleckstrin-homology (PH) region, a C1 sub-type zinc finger (ZF) domain, an SH2 and a C-terminal SH3 (CSH3) (Fig. 1). Additional domains were sequentially acquired into the molecule during evolution, including a proline-rich region (PRR) and a more N-terminally-located SH3 (NSH3) (Fig. 1). The PRR was incorporated in the transition between nematodes and flies. The NSH3 was acquired at the level of tunicates and, since then, remained conserved in all species of the Chordate phyla. The annotation of recently sequenced genomes has revealed that the remnant structure of invertebrate Vav proteins can be found in some Chordate Vav proteins due to differential splicing events. Thus, it is likely that the incorporation of the NSH3 could represent a gain-of-function event not necessarily required for maintaining the basal activity of the full-length proteins. This is consistent with mutagenesis experiments indicating that the functional integrity of the NSH3 domain is not essential for the activation step and effector functions of Vav family proteins in most cell types.<sup>18,19</sup> An important structural property of Vav proteins is that some of their domains form superorder structures in the molecule. For example, the DH, PH, and ZF domains fold together into a common structural unit (referred hereafter as “catalytic core”)<sup>20-22</sup> that, depending on the spatial distribution of the domains, has either positive or negative effects in the overall DH domain catalytic activity.<sup>20</sup> This central cassette also establishes phosphorylation-dependent interactions with the CH-Ac and the CSH3 regions,<sup>20,23</sup> a flip-flop mechanism essential for the control of the overall biological activity of these proteins during signal transduction.<sup>4,18,20,23,24</sup> The Vav PRR and NSH3 domain also form a single functional unit that facilitates the interaction with protein partners.<sup>25,26</sup> Vav proteins are the only Ras superfamily GEFs that harbor a DH-PH-ZF cassette and, consistent with their close connection to tyrosine phosphorylation events, the prototypical


Rho subfamily GTPases, a catalytic step that favors the transition of these GTP-binding proteins from the inactive (GDP bound) to the active, GTP bound state.<sup>4,24,27-29</sup> The main in vitro substrates for Vav proteins include Rac1 and RhoA subfamily proteins.<sup>4,22-24,27-31</sup> Consistent with those catalytic functions, Vav protein can engage a large number of Rho family-dependent cytoskeletal, signaling, and biological responses in most cell types and species tested so far.<sup>12-14</sup> These functions are highly conserved, because active versions of *D. melanogaster* Vav can trigger cytoskeletal change and cell transformation when ectopically expressed in mouse cells.<sup>31</sup> By contrast, Vav proteins do not seem to effectively target Cdc42 under standard catalytic/substrate ratio conditions.<sup>4,23,24,27-30</sup> It should be noted, however, that a very limited number of biological processes regulated by Vav proteins have been attributed, at least partially to Cdc42-dependent mechanisms.<sup>32-42</sup> Whether this is due to catalytic or noncatalytic processes, remains to be determined. Interestingly, in the case of Vav1, the catalysis of nucleotide exchange on Rho proteins requires the cooperation of the DH, PH and ZF regions.<sup>21,22</sup> The requirement of the DH and ZF region seems to be also required in all Vav proteins tested so far. However, there is more flexibility regarding the involvement of the PH region, since constitutively active versions of both Vav2 and Vav3 with inactive PH domains are still catalytically active in vitro (Vav2, Vav3) and in vivo (Vav3).<sup>21,22,43</sup>

The most characteristic noncatalytic function of Vav proteins is the activation of the nuclear factor of activated T-cells (NFAT) in lymphocytes. Although not fully characterized yet, this route requires signals emanating from the Vav CH domain, the Vav-dependent activation of phospholipase C- $\gamma$  (PLC- $\gamma$ ) and the downstream Ca<sup>2+</sup>-dependent calcineurin phosphatase, and synergistic signals triggered by upstream antigenic receptors.<sup>6,8,18,44-48</sup> It probably involves other Vav family- and cell type-specific features as well, because Vav2 and Vav3, unlike the case of Vav1, can trigger NFAT activation in B- but not T-cells.<sup>23,49</sup> Noncatalytic roles for Vav1 in T lymphocytes have been also linked to other biological processes such as integrin-mediated spreading

and the stabilization of Slp-76 microclusters upon T-cell receptor (TCR) stimulation.<sup>50,51</sup> The effectors that mediate such functions are still unknown. The identification of ZF- and CSH3-binding proteins that are not obviously connected to the catalytic performance of Vav1 also suggests that additional catalysis-independent signaling routes might exist<sup>5,6,8,11-14,52</sup> (Fig. 2). Experimental evidence indicates that Vav2 and *D. melanogaster* Vav proteins can also trigger GTPase-independent routes in breast cancer cells and during eye development, respectively<sup>53</sup> (D. Martín-Bermudo, personal communication). Some studies have also described noncatalytic, nuclear roles for Vav proteins. Thus, Vav1 and Vav3 have been associated with transcriptional responses mediated by PU.1, NFAT, NFκB, heat shock transcription factor 1, and the androgen receptor.<sup>54-60</sup> In the case

of Vav3, this transcriptional activity seems to rely on adaptor functions localized within the DH-PH-ZF cassette<sup>57,58,60</sup> (Fig. 2).

Many of the pathways stimulated by cytoplasmic Vav proteins contribute to signaling diversification events via the generation of second messengers such as diacylglycerol (DAG), inositoltriphosphate (IP<sub>3</sub>), Ca<sup>2+</sup>, phosphatidylinositol (3,4,5)-triphosphate (PIP<sub>3</sub>), phosphatidic acid, arachidonic acid, and cyclic guanosine monophosphate (cGMP). The best example for such an activity is seen in lymphocytes, where Vav1 favors the stepwise activation of PLC-γ, phosphatidylinositol-3 kinase (PI3K), protein kinase C (PKC), and PKD.<sup>48,61,62</sup> This leads to the stimulation of non-conventional routes for a Rho GEF such as, for example, the DAG-mediated stimulation of the RasGRP1-Ras-Raf-Erk axis in stimulated T- and B-cells.<sup>8,62-66</sup> Signaling diversification processes have been also found in other cell types such as mouse natural killer cells, mast cells, osteoclasts, vascular smooth muscle cells, and *C. elegans* tissues.<sup>67-71</sup> Depending on the cell type and signaling response involved, Vav proteins can trigger these signaling diversification events in either a catalysis-dependent (i.e., cGMP production in vascular smooth muscle cells, DAG and IP<sub>3</sub> production in B lymphocytes, IP<sub>3</sub> production in *C. elegans* tissues) or independent (i.e., DAG and IP<sub>3</sub> production in T lymphocytes) manner.<sup>47,62,70,71</sup>



**Figure 2.** The Vav family interactome. Examples of some of the Vav family interacting proteins and domains involved. Effector molecules are shown in blue. Positive and negative regulators are shown in green and red, respectively. Proteins whose interaction was only described for Vav2 or Vav3 are indicated by asterisks. Please, note that some of these proteins may exert several functions in the context of Vav signaling.

A still poorly addressed issue is the level of functional redundancy among the 3 Vav family proteins present in Chordate species. However, cell biology and biochemical data suggest that Vav proteins play overlapping, but not identical functions. This evidence includes the different morphologies exhibited by Vav1 and Vav2 oncogene-transformed cells, the synergistic effects of Vav2 and Vav3 in the generation of the Vav2;Vav3-dependent transcriptomal program of breast cancer cells, the Vav1-specific activation of NFAT in T-lymphocytes, the PH requirement for biological activity, and differences in the spectrum of Vav CSH3 binding proteins.<sup>3,4,18,49,53,72-74</sup> It remains to be determined whether those nonredundant activities can be attributed to different subcellular localizations, activities, or spectra of binding partners. To date, nothing is known regarding the signaling specificity of protein isoforms generated by each Vav family gene by differential splicing in some tissues.

## In vivo Roles

Genetic analyses have demonstrated that Vav proteins play key signaling roles in many animal metazoans. In the case of *C. elegans*, the depletion of Vav leads to larval lethality due to

defective rhythmic contractile responses in a number of tissues, including the pharynx, ovaries, and intestine. This phenotype has been associated to Rho- and IP<sub>3</sub>-dependent Ca<sup>2+</sup> generation defects.<sup>71</sup> In *D. melanogaster*, the elimination of Vav leads to defective neuronal axon guidance-related events and embryonic lethality. The few Vav-less flies that manage to hatch show extensive locomotor coordination defects caused by abnormal axon innervation of the ellipsoid body, a key locomotor center in *Diptera*. These defects are due to the normal implication of this protein in Rac1-dependent axon guidance and repulsion processes.<sup>75,76</sup> *D. melanogaster* Vav is also involved in the development of the eye using, as indicated above, catalysis-independent mechanisms (D. Martín-Bermudo, personal communication).

In mammals, the elimination of the haematopoietic-specific Vav1 protein, either alone or in combination with other Vav family proteins, leads to severe defects in lymphocyte development, selection, and effector functions. The signaling dissection of those defects indicate that Vav proteins are essential for T-cell and B-cell receptor responses that contribute to cytoskeletal rearrangement, formation of lymphocyte/antigen presenting cell contacts, PI3K- and PLC- $\gamma$ -mediated signaling diversification events, activation of downstream transcriptional factors, expression of lymphocyte surface markers and cytokines, regulation of cell cycle-promoting proteins, and stimulation of survival pathways. The function of Vav proteins in lymphocytes has been extensively reviewed before.<sup>7,8,12-14</sup> It is likely that Vav1 would play additional roles in T-cells, because the lack of Vav1 unexpectedly leads to the long-term development of T-cell lymphomas, a process further aggravated when the *Rasgrf2* gene is simultaneously knocked out in mice.<sup>77</sup> The mechanistic basis for this “tumor suppressor”-like role remains unknown. Milder effects have been also observed in other haematopoietic cell types, including stem cells, natural killer cells, platelets, neutrophils, macrophages, foam cells, mast cells, and dendritic cells.<sup>12-14</sup>

Outside the haematopoietic system, Vav2 is important for the nitric oxide-mediated relaxation of blood vessels. Due to this, its depletion leads to hypertension and hypertension-associated diseases such as cardiovascular remodeling, cardiorenal fibrosis, and kidney function alterations.<sup>70,78</sup> Vav3 plays roles in bone remodeling, cerebellum development, and axon wiring events in the ventrolateral medulla. The former function is connected to the control of macrophage-colony stimulation factor- and  $\alpha_v\beta_3$  integrin-dependent late osteoclast maturation stages.<sup>69</sup> In the cerebellum, Vav3 is involved in Purkinje cell dendritogenesis, the survival and migration of granule cells, and the formation of the intercrural fissure during the perinatal period. Due to these roles, the absence of Vav3 leads to motor coordination and gaiting defects in early postnatal periods of knockout mice.<sup>79</sup> Finally, Vav3 controls the migration of axons of GABAergic neurons from the caudal to the rostral region of the ventrolateral medulla,<sup>80</sup> a brainstem center that controls sympathetic, respiratory, heart, cardiovascular, and renal activities.<sup>81</sup> When Vav3 is absent, the lack of GABAergic wiring eliminates the tonic inhibition exerted by the caudal over the rostral ventrolateral medulla, leading to chronic sympathoexcitation and a large variety of sympathetic-


mediated defects such as tachycardia, brachypnea, hypertension, and metabolic syndrome.<sup>80,82,83</sup> The analysis of double *Vav2*<sup>-/-</sup>; *Vav3*<sup>-/-</sup> and triple *Vav1*<sup>-/-</sup>; *Vav2*<sup>-/-</sup>; *Vav3*<sup>-/-</sup> knockout mice has revealed defects in axon guidance, dendritic spine formation, and synaptic plasticity in retinal, cortical and hippocampal neurons.<sup>84,85</sup> Alterations in enterocytic differentiation, intestinal epithelial barrier integrity, and angiogenesis have been also observed.<sup>86,87</sup> It is not known as yet whether these functions depend on either the single or combined action of these 3 GEFs.

Genetic data indicate that Vav proteins may cooperate with other Rho GEFs in some biological responses. Those include the collaboration of *D. melanogaster* Vav and Trio during axon guidance processes, the cooperation shown by mouse Vav1 and RasGRF2 proteins in both immature and tumor mouse T-cells, and the functional interaction between Vav1 and P-Rex1 in mouse neutrophil responses.<sup>75,77,88,89</sup> However, despite the multiple Rho GEFs coexpressed with Vav proteins in normal tissues and disease conditions, the inactivation of Vav proteins is sufficient to generate a significant impact in many signaling contexts. This indicates that they regulate quite specific regulatory steps that cannot be redundantly stimulated by other Rho GEFs.

## Regulatory Mechanisms

The biological activity of Vav proteins is controlled by phosphorylation-dependent changes in their intramolecular architecture. Nonphosphorylated Vav proteins are inactive due to a “closed” conformation mediated by extensive intramolecular interactions between the CH-Ac, the CSH3, and the catalytic core (Fig. 3A–C). These contacts occlude the GTPase binding site, favor a catalytically incompetent conformation of the DH-PH-ZF cassette, and limit the noncatalytic output of these proteins.<sup>4,18,20,23,24,30</sup> This inhibited structure is unleashed upon the phosphorylation of Vav proteins on a number of tyrosine residues located in the Ac (Tyr<sup>174</sup> and, to a lower extent Tyr<sup>142</sup>, Tyr<sup>160</sup>), ZF (Tyr<sup>541</sup> and Tyr<sup>544</sup>), and CSH3 (Tyr<sup>836</sup>; amino acid numbers correspond to the primary sequence of mouse Vav1)<sup>20,23,30,90</sup> (Fig. 3A), leading to the activation of both the catalytic and noncatalytic functions of these proteins.<sup>4,18,20,23,24,30,90</sup> The actual structure of this “open” conformation has not been elucidated as yet in the case of the full-length Vav proteins. Due to this, it remains to be determined whether the CH-Ac and CSH3 fully detach from the rest of the domains of the molecule during this activation step or, alternatively, remain in contact with other domains in a different spatial orientation. However, structural studies conducted with N-terminally truncated Vav1 indicate that the catalytic DH-PH-ZF core of Vav proteins probably undergoes a shift from a catalytically incompetent to a catalytically competent conformation upon the release of the inhibitory domains.<sup>20</sup> Likewise, the CSH3 seems to remain in contact with the catalytic core when Vav proteins are in the “open” state.<sup>23</sup> In addition to the exposure of effector sites, it is possible that this flip-flop activation could be used for other regulatory layers. For example, it can be speculated that the need of multiple phosphorylation events for






optimal Vav activation could be used to establish minimal, digital-like thresholds for the activation of these proteins, setting up gradients of activity, or engaging different downstream routes depending on the strength of incoming extracellular stimuli and the total number of phosphosites engaged at a given time. Conversely, they can represent a safeguard mechanism to avoid the spurious stimulation of Vav proteins by the stochastic phosphorylation of single phosphosites in nonstimulated cells. Interactions of the Vav CSH3 with other proteins in the “close” state may also induce conformational changes that could facilitate the access of the hidden tyrosine hydroxyl groups for the phosphorylation step. Alternatively, they could delay the return of the CSH3 back to the “close” state when taking place in the active conformation, thus facilitating either the time- or subcellular location-specific extension of active Vav signaling output in cells. Regardless of all its potential regulatory roles, the relevance of this intramolecular mechanism is underscored by its phylogenetic conservation and, in addition, by the hyperactivation observed in Vav proteins upon the mutation of either domains or residues involved in maintaining the autoinhibited conformation<sup>3,4,18,20,23,24,31,71,90</sup> (Fig. 3D and E). Indeed, a 66 amino acid-long deletion of the Vav1 CH domain was the genetic alteration that allowed the initial discovery of the *VAV1* oncogene<sup>1,91</sup> (Fig. 3D). It is worth noting that phosphorylation-independent activation steps for the Vav family have been observed upon the interaction of the p67 phagocyte oxidase subunit (p67<sup>Phox</sup>) and the human immunodeficiency virus (HIV)-encoded v-Nef protein with the Vav1 catalytic core and CSH3, respectively<sup>41,92</sup> (Fig. 2). The basis for such activation remains unknown, although it may involve domain displacement mechanisms that could favor the spurious “opening” of the autoinhibited state of Vav proteins. This type of domain displacement-based activation has been found before in other signaling molecules, such as the Src kinase family.<sup>93</sup>

The Vav phosphorylation step is mediated by either transmembrane or cytosolic PTKs. The former subclass encompasses a large variety of receptors for both soluble growth factors and ephrins. The phosphorylation of Vav proteins by these receptors usually involves a direct, SH2-mediated interaction of Vav proteins with the autophosphorylated cytoplasmic tails of the kinases.<sup>5,12-14</sup> Historically, this was actually the first connection ever described between Vav proteins and PTKs.<sup>94,95</sup> Cytosolic Src, Syk, Janus, Tec, and Abl family kinases are mainly involved in the phosphorylation of Vav proteins downstream of receptors lacking intrinsic tyrosine kinase domains such as those for antigens, cytokines, and extracellular matrix proteins (i.e., integrins).<sup>5,12-14</sup> Src kinases are also in charge of phosphorylating Vav proteins in other signaling cascades, including those initiated by nitric oxide, Wnt, tumor necrosis factor, diacylglycerol-regulated protein kinases, and some transmembrane PTKs.<sup>70,96-99</sup> The close connection of Src PTKs with Vav proteins probably extends far in evolution, because these molecules are one of the few kinase families present in Choanoflagellates.<sup>15</sup> Although the interaction of Vav proteins with cytosolic kinases may be direct as in the case of the transmembrane PTKs (Fig. 4, point a), it usually entails the prior orchestration of either 2 or 3-step plasma

membrane translocation mechanisms. In the former case, Vav proteins establish SH2-mediated interactions with phosphorylated coreceptors (i.e., CD19, Lck interacting membrane protein)<sup>49,100-102</sup> (Fig. 4, point b). In the latter case, Vav proteins first interact with cytosolic adaptor molecules using the SH2 (Slp76, Blnk), PRR-NSH3 (Grb2, Nck) or the CSH3 (dynamin 2) domains<sup>51,74,103-109</sup> and, subsequently, utilize those adaptors to dock onto phosphorylated proteins localized at the plasma membrane (Fig. 4, point c). These docking proteins include coreceptors (CD28), membrane-anchored scaffolding proteins (i.e., Lat), and TCR-localized adaptor molecules (Shb).<sup>106,110,111</sup> Regardless of the translocation mechanism involved, the endpoint result is the phosphorylation of Vav proteins by the upstream cytoplasmic PTKs. This latter step involves transient physical interactions with the kinase, since the Vav SH2 domains are always essential for the effective phosphorylation of these proteins.<sup>18,19</sup> It is expected that Vav proteins will return to the inactive, “close” state at the end of the stimulation cycle via tyrosine dephosphorylation (see above, Fig. 3A). However, this regulatory step remains poorly characterized.

The translocation of Vav1 to the nucleus depends on a nuclear localization signal present in the PH region and, in addition, by uncharacterized cytoplasmic sequestering functions of the CSH3.<sup>55</sup> However, the mechanism that mediates the translocation step and subsequent activity of Vav1 in the nucleus still remains obscure. Posttranslational modifications could be involved, as inferred from the nucleus-specific arginine methylation of Vav1.<sup>112</sup> The cotranscriptional activity of Vav3 is regulated by the PH-mediated interaction with the cell division cycle 37 cochaperone<sup>60</sup> (Fig. 2).

Vav proteins can be potentially controlled by lysine acetylation, caspase-mediated cleavage, proteolysis, microRNAs, gene expression regulation, and epigenetic mechanisms.<sup>33,113-122</sup> The functional significance of most of these regulatory steps is still unknown. However, there is strong genetic evidence in favor of the microRNA-dependent regulation *C. elegans* Vav during Notch signaling.<sup>117</sup> The concerted action of demethylation and transcriptional factors (i.e., c-Myb) has been shown to be involved in the ectopic expression of human *VAV1* gene in non-hematopoietic cancer cells.<sup>120,122</sup> The mouse *Vav3* gene is also transcriptionally activated by the aryl hydrocarbon receptor (Ahr) under normal physiological conditions.<sup>121,123</sup> Early studies showed that the catalytic activity of Vav proteins could be activated upon the binding of DAG and PIP<sub>3</sub> to the ZF and PH domains, respectively.<sup>124,125</sup> However, such regulatory action seems at odds with the known inability of those domains to directly bind those second messengers.<sup>126-129</sup> Consistent with this, it has been shown that these 2 Vav domains do not undergo the expected DAG- and PIP<sub>3</sub>-dependent changes in subcellular localization when expressed in cells.<sup>127,128</sup> In the case of the ZF, the reason for such lack of binding has been comprehensively analyzed at the structural and biochemical level.<sup>126,127</sup> In the case of the PH, the lack of PIP<sub>3</sub> binding is probably due to the absence of consensus amino acid sequences that typically mediate the high affinity interaction of other PH domains with this second messenger.<sup>128,129</sup> Extensive signaling and genetic evidence



**Figure 4.** Signaling steps for the stimulation of Vav proteins in lymphocytes. For simplicity, we only show the phosphorylation step mediated by a Syk family member. Gray and green arrows represent translocation and phosphorylation steps, respectively. This step should include SH2-mediated interactions with the kinase (not shown in steps b and c). ITAM, immunoreceptor tyrosine-based activation motif.

also indicate that Vav proteins act upstream rather than downstream of both PLC- $\gamma$  and PI3K family proteins.<sup>8,47,62,130</sup>

## Pathological Roles

There are no consistent examples of either gain- or loss-of-function mutations for *VAV* family genes in human disease. However, the regulatory cycle of Vav proteins can be altered in some diseases by increased abundance or upon ectopic expression in tissues that do not normally express them.<sup>53,120,122</sup> In other cases, Vav proteins can become hyperphosphorylated due to the presence of oncogenic PTKs, mitogen-driven autocrine loops, increased abundance of Vav-binding adaptors, and the CSH3-mediated interaction with the  $\gamma$ -herpesvirus M2 adaptor protein.<sup>35,73,131-136</sup> Alternatively, the activation step can take place without apparent changes in the phosphorylation status of Vav proteins, as previously discussed in the case of the HIV v-Nef

protein.<sup>41</sup> Genomic studies have revealed that single nucleotide polymorphisms (SNPs) in *VAV* genes seem to be linked to multiple sclerosis (*VAV1*), schizophrenia (*VAV3*), and hypothyroidism (*VAV3*).<sup>137-139</sup> It is still unknown whether there is a cause-effect relationship between most of those genetic linkages and the evolution of those diseases. The only exception is multiple sclerosis, since patients containing specific SNPs in the first intron of the *VAV1* gene show elevated abundance of the *VAV1* mRNA and, in addition, increased production of proinflammatory cytokines by leukocytes present in both peripheral blood and the cerebrospinal fluid.<sup>137</sup> Conversely, the presence of a specific SNP in the first exon of the rat *Vav1* gene that promotes the expression of a poorly expressed Vav1 mutant version (R63W) leads to reduced production of proinflammatory factors and increased resistance to autoimmune encephalitis in this rodent species.<sup>137</sup> Specific *VAV2* and *VAV3*

SNPs have been also associated with glaucoma development,<sup>140</sup> although this linkage has not been corroborated by independent studies using other patient cohorts.<sup>141,142</sup>

Experiments with knockout mice indicate that endogenous Vav proteins do play important roles in the evolution and/or progression of some of the above diseases. For example, the Vav1-Rac1;Rac2-Pak1 pathway has been shown to be important for the development and maintenance of c-Kit-positive acute myeloid leukemia through a cancer cell survival mechanism.<sup>143</sup> Vav3 is important for the fitness of p190<sup>Bcr-Abl</sup>-driven acute lymphoblastic leukemia, an effect attributed to the implication of this family member in the Rac2- and Pak1-dependent regulation of both pro- and antiapoptotic Bcl family members.<sup>144</sup> Vav2 and Vav3 proteins are both required for breast tumorigenesis as well as for the initiation and promotion phases of skin tumors.<sup>53,97</sup> In the case of breast, these proteins promote distal transcriptional programs affecting the proliferation, survival, and neoangiogenesis of cells in the primary tumor. In addition, they favor the



extravasation and subsequent fitness of breast cancer cells in the lung parenchyma.<sup>53</sup> In the skin, they make it possible the expression of large autocrine and paracrine programs that facilitate the keratinocyte survival to DNA damaging insults, keratinocyte G<sub>1</sub>-S cell cycle transitions during the tumor promotion phase, and the generation of a proinflammatory environment.<sup>97</sup> Interestingly, the lack of Vav2 and Vav3 does not induce any overt defects in the normal development, function, and long-term homeostasis of both mammary glands and skin.<sup>53,97</sup> This suggests that Vav proteins may exert disease-specific tasks not directly connected to their normal function in healthy tissues. The contribution of Vav proteins to some of those tumors seems to be seminal, as inferred from the detection of Vav2;Vav3-dependent breast cancer gene signatures that can predict disease outcome in breast cancer patients.<sup>53</sup>

Outside the cancer field, the systemic elimination of Vav1 blocks hyperlipidemia-linked prothrombotic states as well formation of atherosclerotic lesions. These therapeutic effects are due to the specific function of Vav1 in platelets and macrophage-derived foam cells, respectively.<sup>145-147</sup> The specific elimination of Vav1 catalytic activity has been shown to be sufficient for eliminating other clinically relevant problems such as allograft tissue rejection and graft versus host disease.<sup>148</sup> Unlike the case of breast and skin tumors, these latter results probably reflect the intrinsic role of Vav1 in normal T-lymphocyte signaling. Vav2 is also important for the development of hyperhomocysteinemia-induced renal glomerulosclerosis,<sup>149</sup> a pathophysiological event commonly associated to end-stage renal disease in hypertensive patients. This effect has been linked to defective reactive oxygen species production in the kidney during those conditions.<sup>149</sup> Finally, a number of genetic and cell biology approaches have demonstrated the implication of Vav proteins, namely Vav1, in the pathogenic cycles of both HIV and  $\gamma$ -herpesviruses.<sup>41,73</sup>

## Therapeutic Value of Vav Proteins: Pros and Cons

The above data indicate that Vav proteins could represent potentially interesting therapeutic targets. However, this is far from being an open and shut case. Thus, a devil's advocate would probably dismiss these knockout-based data because they do not formally prove that the inactivation of Vav proteins is effective in already developed diseases and, even if that were the case, whether the simple inhibition of their catalytic activities would be sufficient to generate a noticeable therapeutic impact. To tackle this issue, it will be important to generate "second generation" knock-in mice capable of expressing catalytically inactive Vav proteins in an inducible manner at clinically relevant disease progression stages. The devil's advocate could also argue that the collateral effects induced by the inhibition of Vav proteins in healthy tissues might hamper the implementation of such anti-Vav therapies. According to the data generated with Vav family knockout mice, such side effects could include hypertension, metabolic syndrome, infection-associated endotoxemia, and immunosuppression.<sup>8,54,78,82,83</sup> Some of these problems stem from roles of Vav3 in the embryonic period or, in the case of

endotoxemia, in noncatalytic functions of Vav1.<sup>54,80</sup> It is predictable therefore that they will not show when using catalytic inhibitors for Vav proteins in adult patients. However, the inactivation of Vav2 in postnatal periods should promote hypertension and associated diseases.<sup>70</sup> This will actually represent a common problem for most Rac1-based therapies, because the inducible smooth muscle cell-specific inactivation of the Rac1 gene in adult mice triggers the same defects.<sup>150,151</sup> However, these side effects can be prevented, or cured if already developed, using standard antihypertensive treatments currently available in the clinic<sup>70,78,82</sup> (S. Fabbiano and X.R.B., manuscript submitted). The systemic inhibition of Vav proteins will also lead to immunosuppression and, possibly, minor defects in other haematopoietic lineages.<sup>8,12-14</sup> However, we do not know yet whether such defects would arise upon the inducible inhibition of Vav1 or, alternatively, in adulthood periods when thymi have already regressed. It is also possible that, as in many other targeted therapies, clinical benefits could be obtained without dire side effects due to the high signaling and metabolic demands for cells in pathological settings. Evidence for such therapeutic windows does exist in the case of Vav2 and Vav3 in both the mammary gland and skin.<sup>53,97</sup>

Another important issue is whether we could successfully isolate pharmacologically effective chemical inhibitors for these proteins. Indeed, it is widely assumed that the isolation of high affinity inhibitors to the catalytic site of Rho GEFs is rather difficult due to the shallow nature of the GEF/GTPase binding interface.<sup>152</sup> However, the idiosyncratic nature of the catalytic core of Vav proteins and its mechanism of activation can offer alternative ways to achieve such inhibition without touching the catalytic site. For example, the DH-PH-ZF structure contains additional pockets for effective inhibition, a strategy that has been proven fruitful before in the case of other Rho GEF interdomain linker regions.<sup>153</sup> Alternatively, we could generate drugs that could mimic the inhibitory CH-Ac and CSH3 interactions with the catalytic core. Finally, Vav-dependent diseases could be probably treated using indirect approaches, such as either the single or combined inhibition of therapeutic targets encoded by Vav-regulated genes. Experimental evidence supporting this alternative approach has been recently obtained in Vav-dependent breast cancer cells.<sup>53</sup> In any case, and as discussed above, the success of this type of therapies will ultimately depend on the relative contribution of the catalytic function of these proteins to the fitness of already developed diseases. Although intrinsic to any type of target-directed therapies, all the aforementioned problems underscore the need of deepening our understanding of the role of Vav proteins, and especially of their catalysis-dependent pathways, in both normal physiological and disease conditions.

## Final Remarks

Despite significant advances made in the understanding of the regulatory and effector properties of Vav proteins, there are still significant open questions in this field. Thus, we have still to get a holistic view of the regulatory landscape of these proteins, the contribution of GTPase-dependent and independent routes to



the overall biological program of each Vav protein, and the level of overlap/signaling specificity that exists among them. A better understanding of Vav2 and Vav3 is still required because, up to now, most of the characterization of the functional properties of this family has been Vav1-based. Likewise, the function of Vav proteins in many model organisms is still unknown. We also need to complete the spectrum of Vav-dependent diseases and the minimal number of Vav proteins that have to be inhibited to get optimal therapeutic effects. In this context, it would be of paramount importance to generate new animal models to assess the therapeutic value of targeting the catalytic activity of these proteins and, in addition, the side effects derived from such inhibition. Finally, we have to devise ways to generate high affinity, Vav family-specific inhibitors. This future work will give us a holistic understanding of these proteins and, perhaps more importantly, yield innovative tools to both diagnose and treat human diseases.

## References

- Katzav S, Martin-Zanca D, Barbacid M. Vav, a novel human oncogene derived from a locus ubiquitously expressed in hematopoietic cells. *EMBO J* 1989; 8:2283-90; PMID:2477241
- Henske EP, Short MP, Jozwiak S, Bovey CM, Ramla-khan S, Haines JL, Kwiatkowski DJ. Identification of VAV2 on 9q34 and its exclusion as the tuberous sclerosis gene TSC1. *Ann Hum Genet* 1995; 59:25-37; PMID:7762982; <http://dx.doi.org/10.1111/j.1469-1809.1995.tb01603.x>
- Schubel KE, Bustelo XR, Nielsen DA, Song BJ, Barbacid M, Goldman D, Lee IJ. Isolation and characterization of murine vav2, a member of the vav family of proto-oncogenes. *Oncogene* 1996; 13:363-71; PMID: 8710375
- Movilla N, Bustelo XR. Biological and regulatory properties of Vav-3, a new member of the Vav family of oncoproteins. *Mol Cell Biol* 1999; 19:7870-85; PMID:10523675
- Bustelo XR. Regulatory and signaling properties of the Vav family. *Mol Cell Biol* 2000; 20:1461-77; PMID:10669724; <http://dx.doi.org/10.1128/MCB.20.5.1461-1477.2000>
- Bustelo XR. Vav proteins, adaptors and cell signaling. *Oncogene* 2001; 20:6372-81; PMID:11607839; <http://dx.doi.org/10.1038/sj.onc.1204780>
- Turner M, Billadeau DD. VAV proteins as signal integrators for multi-subunit immune-recognition receptors. *Nat Rev Immunol* 2002; 2:476-86; PMID:12094222; <http://dx.doi.org/10.1038/nri840>
- Tybulewicz VL. Vav-family proteins in T-cell signalling. *Curr Opin Immunol* 2005; 17:267-74; PMID: 15886116; <http://dx.doi.org/10.1016/j.coi.2005.04.003>
- Bustelo XR. Regulation of Vav proteins by intramolecular events. *Front Biosci* 2002; 7:d24-30; PMID:11779690; <http://dx.doi.org/10.2741/bustelo>
- Hornstein I, Alcover A, Katzav S. Vav proteins, masters of the world of cytoskeleton organization. *Cell Sig* 2004; 16:1-11; PMID:14607270; [http://dx.doi.org/10.1016/S0898-6568\(03\)00110-4](http://dx.doi.org/10.1016/S0898-6568(03)00110-4)
- Bustelo XR. Vav family. In: Choy S, ed. *Encyclopedia of Signaling Molecules*. New York: Springer, 2012; 1963-76
- Bustelo XR. Vav1. *UCSD Nat Mol Pages* 2010; <http://dx.doi.org/10.1038/mp.a002360.01>
- Bustelo XR, Couceiro JR. Vav3. *UCSD Nat Mol Pages* 2008; <http://dx.doi.org/10.1038/mp.a002362.01>
- Bustelo XR. Vav2. *UCSD Nat Mol Pages* 2008; <http://dx.doi.org/10.1038/mp.a002361.01>
- Manning G, Young SL, Miller WT, Zhai Y. The protist, *Monosiga brevicollis*, has a tyrosine kinase signaling network more elaborate and diverse than found in

- any known metazoan. *Proc Natl Acad Sci USA* 2008; 105:9674-9; PMID:18621719; <http://dx.doi.org/10.1073/pnas.0801314105>
- Bustelo XR, Rubin SD, Suen KL, Carrasco D, Barbacid M. Developmental expression of the vav proto-oncogene. *Cell Growth Differ* 1993; 4:297-308; PMID:8494792
- Trenkle T, McClelland M, Adlkofer K, Welsh J. Major transcript variants of VAV3, a new member of the VAV family of guanine nucleotide exchange factors. *Gene* 2000; 245:139-49; PMID:10713454; [http://dx.doi.org/10.1016/S0378-1119\(00\)00026-3](http://dx.doi.org/10.1016/S0378-1119(00)00026-3)
- Zugaza JL, Lopez-Lago MA, Caloca MJ, Dosil M, Movilla N, Bustelo XR. Structural determinants for the biological activity of Vav proteins. *J Biol Chem* 2002; 277:45377-92; PMID:12228230; <http://dx.doi.org/10.1074/jbc.M208039200>
- Ksionda O, Saveliev A, Kochl R, Rapley J, Faroudi M, Smith-Garvin JE, Wulffing C, Rittinger K, Carter T, Tybulewicz VL. Mechanism and function of Vav1 localisation in TCR signalling. *J Cell Sci* 2012; 125:5302-14; PMID:22956543; <http://dx.doi.org/10.1242/jcs.105148>
- Yu B, Martins IR, Li P, Amarasinghe GK, Umetani J, Fernandez-Zapico ME, Billadeau DD, Machius M, Tomchick DR, Rosen MK. Structural and energetic mechanisms of cooperative autoinhibition and activation of Vav1. *Cell* 2010; 140:246-56; PMID: 20141838; <http://dx.doi.org/10.1016/j.cell.2009.12.033>
- Rapley J, Tybulewicz VL, Rittinger K. Crucial structural role for the PH and C1 domains of the Vav1 exchange factor. *EMBO Rep* 2008; 9:655-61; PMID: 18511940; <http://dx.doi.org/10.1038/embor.2008.80>
- Chrencik JE, Brooun A, Zhang H, Mathews II, Hura GL, Foster SA, Perry JJ, Streiff M, Ramage P, Widmer H, et al. Structural basis of guanine nucleotide exchange mediated by the T-cell essential Vav1. *J Mol Biol* 2008; 380:828-43; PMID:18589439; <http://dx.doi.org/10.1016/j.jmb.2008.05.024>
- Barreira M, Fabbiano S, Couceiro JR, Torreira E, Martinez-Torrecuadrada JL, Montoya G, Llorca O, Bustelo XR. The C-terminal SH3 domain contributes to the intramolecular inhibition of Vav family proteins. *Sci Sig* 2014; 7:ra35; PMID:24736456; <http://dx.doi.org/10.1126/scisignal.2004993>
- Schubel KE, Movilla N, Rosa JL, Bustelo XR. Phosphorylation-dependent and constitutive activation of Rho proteins by wild-type and oncogenic Vav-2. *EMBO J* 1998; 17:6608-21; PMID:9822605; <http://dx.doi.org/10.1093/emboj/17.22.6608>
- Nishida M, Nagata K, Hachimori Y, Horiuchi M, Ogura K, Mandiyan V, Schlesinger J, Inagaki F. Novel recognition mode between Vav and Grb2 SH3

- domains. *EMBO J* 2001; 20:2995-3007; PMID: 11406576; <http://dx.doi.org/10.1093/emboj/20.12.2995>
- Ogura K, Nagata K, Horiuchi M, Ebisui E, Hasuda T, Yuzawa S, Nishida M, Hatanaka H, Inagaki F. Solution structure of N-terminal SH3 domain of Vav and the recognition site for Grb2 C-terminal SH3 domain. *J Biomol NMR* 2002; 22:37-46; PMID: 11885979; <http://dx.doi.org/10.1023/A:1013868731495>
- Crespo P, Schubel KE, Ostrom AA, Gutkind JS, Bustelo XR. Phosphotyrosine-dependent activation of Rac-1 GTPGTP exchange by the vav proto-oncogene product. *Nature* 1997; 385:169-72; PMID:8990121; <http://dx.doi.org/10.1038/385169a0>
- Movilla N, Dosil M, Zheng Y, Bustelo XR. How Vav proteins discriminate the GTPases Rac1 and RhoA from Cdc42. *Oncogene* 2001; 20:8057-65; PMID: 11781818; <http://dx.doi.org/10.1038/sj.onc.1205000>
- Abe K, Rossman KL, Liu B, Ritola KD, Chiang D, Campbell SL, Burrig K, Der CJ. Vav2 is an activator of Cdc42, Rac1, and RhoA. *J Biol Chem* 2000; 275:10141-9; PMID:10744696; <http://dx.doi.org/10.1074/jbc.275.14.10141>
- Aghazadeh B, Lowry WE, Huang XY, Rosen MK. Structural basis for relief of autoinhibition of the Dbl homology domain of proto-oncogene Vav by tyrosine phosphorylation. *Cell* 2000; 102:625-33; PMID: 11007481; [http://dx.doi.org/10.1016/S0092-8674\(00\)00085-4](http://dx.doi.org/10.1016/S0092-8674(00)00085-4)
- Couceiro JR, Martin-Bermudo MD, Bustelo XR. Phylogenetic conservation of the regulatory and functional properties of the Vav oncoprotein family. *Exp Cell Res* 2005; 308:364-80; PMID:15950967; <http://dx.doi.org/10.1016/j.yexcr.2005.04.035>
- Razidlo GL, Schroeder B, Chen J, Billadeau DD, McNiven MA. Vav1 as a central regulator of invadopodia assembly. *Curr Biol* 2014; 24:86-93; PMID: 24332539; <http://dx.doi.org/10.1016/j.cub.2013.11.013>
- Duan L, Raja SM, Chen G, Virmani S, Williams SH, Clubb RJ, Mukhopadhyay C, Rainey MA, Ying G, Dimri M, et al. Negative regulation of EGFR-Vav2 signaling axis by Cbl ubiquitin ligase controls EGF receptor-mediated epithelial cell adherens junction dynamics and cell migration. *J Biol Chem* 2011; 286:620-33; PMID:20940296; <http://dx.doi.org/10.1074/jbc.M110.188086>
- Duan L, Chen G, Virmani S, Ying G, Raja SM, Chung BM, Rainey MA, Dimri M, Ortega-Cava CF, Zhao X, et al. Distinct roles for Rho versus RacCdc42 GTPases downstream of Vav2 in regulating mammary epithelial acinar architecture. *J Biol Chem* 2010; 285:1555-68; PMID:19826000; <http://dx.doi.org/10.1074/jbc.M109.057976>

## Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

## Funding

Our Vav-related work has been funded by the Spanish Ministry of Economy and Competitiveness (SAF2012-3171, RD12/0036/0002), the Castilla-León Autonomous Government (CSI101U13), the US National Institutes of Health-National Cancer Institute (RO1CA7373509), the European Union 7<sup>th</sup> Framework Program (FP7-HEALTH-2007-A-201862), Worldwide Cancer Research (14-1248), the Asociación Española contra el Cáncer, and the Ramón Areces Foundation. Spanish funding is cosponsored by the European Regional Development Fund. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

35. Ambrogio C, Voena C, Manazza AD, Martinengo C, Costa C, Kirchhausen T, Hirsch E, Inghirami G, Chiarle R. The anaplastic lymphoma kinase controls cell shape and growth of anaplastic large cell lymphoma through Cdc42 activation. *Cancer Res* 2008; 68:8899-907; PMID:18974134 <http://dx.doi.org/10.1158/0008-5472.CAN-08-2568>
36. Aoki K, Nakamura T, Fujikawa K, Matsuda M. Local phosphatidylinositol 3,4,5-trisphosphate accumulation recruits Vav2 and Vav3 to activate Rac1Cdc42 and initiate neurite outgrowth in nerve growth factor-stimulated PC12 cells. *Mol Biol Cell* 2005; 16:2207-17; PMID:15728722; <http://dx.doi.org/10.1091/mbc.E04-10-0904>
37. Tu S, Wu WJ, Wang J, Cerione RA. Epidermal growth factor-dependent regulation of Cdc42 is mediated by the Src tyrosine kinase. *J Biol Chem* 2003; 278:49293-300; PMID:14506284; <http://dx.doi.org/10.1074/jbc.M307021200>
38. Gakidis MA, Cullere X, Olson T, Wilsbacher JL, Zhang B, Moores SL, Ley K, Swat W, Mayadas T, Brugge JS. Vav GEFs are required for beta2 integrin-dependent functions of neutrophils. *J Cell Biol* 2004; 166:273-82; PMID:15249579; <http://dx.doi.org/10.1083/jcb.200404166>
39. Zeng R, Cannon JL, Abraham RT, Way M, Billadeau DD, Bubeck-Wardenberg J, Burkhardt JK. SLP-76 coordinates Nck-dependent Wiskott-Aldrich syndrome protein recruitment with Vav-1Cdc42-dependent Wiskott-Aldrich syndrome protein activation at the T cell-APC contact site. *J Immunol* 2003; 171:1360-8; PMID:12874226; <http://dx.doi.org/10.4049/jimmunol.171.3.1360>
40. Ardouin L, Bracke M, Mathiot A, Pagakis SN, Norton T, Hogg N, Tybulewicz VL. Vav1 transduces TCR signals required for LFA-1 function and cell polarization at the immunological synapse. *Eur J Immunol* 2003; 33:790-7; PMID:12616499; <http://dx.doi.org/10.1002/eji.200323858>
41. Fackler OT, Luo W, Geyer M, Alberts AS, Peterlin BM. Activation of Vav by Nef induces cytoskeletal rearrangements and downstream effector functions. *Mol Cell* 1999; 3:729-39; PMID:10394361; [http://dx.doi.org/10.1016/S1097-2765\(01\)80005-8](http://dx.doi.org/10.1016/S1097-2765(01)80005-8)
42. Liu BP, Burrige K. Vav2 activates Rac1, Cdc42, and RhoA downstream from growth factor receptors but not beta1 integrins. *Mol Cell Biol* 2000; 20:7160-9; PMID:10982832; <http://dx.doi.org/10.1128/MCB.20.19.7160-7169.2000>
43. Booden MA, Campbell SL, Der CJ. Critical but distinct roles for the pleckstrin homology and cysteine-rich domains as positive modulators of Vav2 signaling and transformation. *Mol Cell Biol* 2002; 22:2487-97; PMID:11909943; <http://dx.doi.org/10.1128/MCB.22.8.2487-2497.2002>
44. Wu J, Katav S, Weiss A. A functional T-cell receptor signaling pathway is required for p95vav activity. *Mol Cell Biol* 1995; 15:4337-46; PMID:7623828
45. Kuhne MR, Ku G, Weiss A. A guanine nucleotide exchange factor-independent function of Vav1 in transcriptional activation. *J Biol Chem* 2000; 275:2185-90; PMID:10636924; <http://dx.doi.org/10.1074/jbc.275.3.2185>
46. Zhou Z, Yin J, Dou Z, Tang J, Zhang C, Cao Y. The calponin homology domain of Vav1 associates with calmodulin and is prerequisite to T cell antigen receptor-induced calcium release in Jurkat T lymphocytes. *J Biol Chem* 2007; 282:23737-44; PMID:17550897; <http://dx.doi.org/10.1074/jbc.M702975200>
47. Saveliev A, Vanes L, Ksionda O, Rapley J, Smerdon SJ, Rittinger K, Tybulewicz VL. Function of the nucleotide exchange activity of Vav1 in T cell development and activation. *Sci Sig* 2009; 2:ra83; PMID:20009105; <http://dx.doi.org/10.1126/scisignal.2000420>
48. Reynolds LF, Smyth LA, Norton T, Freshney N, Downward J, Kioussis D, Tybulewicz VL. Vav1 transduces T cell receptor signals to the activation of phospholipase C-gamma1 via phosphoinositide 3-kinase-dependent and -independent pathways. *J Exp Med* 2002; 195:1103-14; PMID:11994416; <http://dx.doi.org/10.1084/jem.200111663>
49. Doody GM, Billadeau DD, Clayton E, Hutchings A, Berland R, McAdam S, Leibson PJ, Turner M. Vav-2 controls NFAT-dependent transcription in B- but not T-lymphocytes. *EMBO J* 2000; 19:6173-84; PMID:11080163; <http://dx.doi.org/10.1093/emboj/19.22.6173>
50. del Pozo MA, Schwartz MA, Hu J, Kiosses WB, Altman A, Villalba M. Guanine exchange-dependent and -independent effects of Vav1 on integrin-induced T cell spreading. *J Immunol* 2003; 170:41-7; PMID:12496381; <http://dx.doi.org/10.4049/jimmunol.170.1.41>
51. Sylvain NR, Nguyen K, Bunnell SC. Vav1-mediated scaffolding interactions stabilize SLP-76 microclusters and contribute to antigen-dependent T cell responses. *Sci Sig* 2011; 4:ra14; PMID:21386095; <http://dx.doi.org/10.1126/scisignal.2001178>
52. Jacob CO, Eisenstein M, Dinauer MC, Ming W, Liu Q, John S, Quismorio FP Jr, Reiff A, Myones BL, Kaufman KM, et al. Lupus-associated causal mutation in neutrophil cytosolic factor 2 (NCF2) brings unique insights to the structure and function of NADPH oxidase. *Proc Natl Acad Sci USA* 2012; 109:E59-67; PMID:22203994; <http://dx.doi.org/10.1073/pnas.1113251108>
53. Citterio C, Menacho-Marquez M, Garcia-Escudero R, Larive RM, Barreiro O, Sanchez-Madrid F, Paramio JM, Bustelo XR. The Rho exchange factors Vav2 and Vav3 control a lung metastasis-specific transcriptional program in breast cancer cells. *Sci Sig* 2012; 5:ra71; PMID:23033540; <http://dx.doi.org/10.1126/scisignal.2002962>
54. Zenker S, Pantelev-Ivlev J, Wirtz S, Kishimoto T, Waldner MJ, Ksionda O, Tybulewicz VL, Neuhart MF, Atreya I. A key regulatory role for Vav1 in controlling lipopolysaccharide endotoxemia via macrophage-derived IL-6. *J Immunol* 2014; 192:2830-6; PMID:24532586; <http://dx.doi.org/10.4049/jimmunol.1300157>
55. Houliard M, Arudchandran R, Regnier-Ricard F, Germani A, Gisselbrecht S, Blank U, Rivera J, Varin-Blank N. Vav1 is a component of transcriptionally active complexes. *J Exp Med* 2002; 195:1115-27; PMID:11994417; <http://dx.doi.org/10.1084/jem.20011701>
56. Brugnoli F, Lambertini E, Varin-Blank N, Piva R, Marchisio M, Grassilli S, Miscia S, Capitani S, Bertagnolo V. Vav1 and PU.1 are recruited to the CD11b promoter in APL-derived promyelocytes: role of Vav1 in modulating PU.1-containing complexes during ATRA-induced differentiation. *Exp Cell Res* 2010; 316:38-47; PMID:19747912; <http://dx.doi.org/10.1016/j.yexcr.2009.09.004>
57. Lyons LS, Burnstein KL. Vav3, a Rho GTPase guanine nucleotide exchange factor, increases during progression to androgen independence in prostate cancer cells and potentiates androgen receptor transcriptional activity. *Mol Endocrinol* 2006; 20:1061-72; PMID:16384856; <http://dx.doi.org/10.1210/me.2005-0346>
58. Liu Y, Wu X, Dong Z, Lu S. The molecular mechanism of Vav3 oncogene on upregulation of androgen receptor activity in prostate cancer cells. *Int J Oncol* 2010; 36:623-33; PMID:20126983; <http://dx.doi.org/10.3892/ijo.00000644>
59. Peacock SO, Fahrenholtz CD, Burnstein KL. Vav3 enhances androgen receptor splice variant activity and is critical for castration-resistant prostate cancer growth and survival. *Mol Endocrinol* 2012; 26:1967-79; PMID:23023561; <http://dx.doi.org/10.1210/me.2012-1165>
60. Wu F, Peacock SO, Rao S, Lemmon SK, Burnstein KL. Novel interaction between the co-chaperone Cdc37 and Rho GTPase exchange factor Vav3 promotes androgen receptor activity and prostate cancer growth. *J Biol Chem* 2013; 288:5463-74; PMID:23281476; <http://dx.doi.org/10.1074/jbc.M112.390963>
61. Vigorito E, Kovess D, Turner M. Synergistic activation of PKD by the B cell antigen receptor and CD19 requires PI3K, Vav1 and PLCgamma. *Cell Sig* 2006; 18:1455-60; PMID:16380231; <http://dx.doi.org/10.1016/j.celsig.2005.11.008>
62. Caloca MJ, Zugaza JL, Bustelo XR. Mechanistic analysis of the amplification and diversification events induced by Vav proteins in B-lymphocytes. *J Biol Chem* 2008; 283:36454-64; PMID:18974050; <http://dx.doi.org/10.1074/jbc.M803814200>
63. Caloca MJ, Zugaza JL, Matallanas D, Crespo P, Bustelo XR. Vav mediates Ras stimulation by direct activation of the GTPGTP exchange factor Ras GRP1. *EMBO J* 2003; 22:3326-36; PMID:12839994; <http://dx.doi.org/10.1093/emboj/cdg316>
64. Zugaza JL, Caloca MJ, Bustelo XR. Inverted signaling hierarchy between RAS and RAC in T-lymphocytes. *Oncogene* 2004; 23:5823-33; PMID:15184873; <http://dx.doi.org/10.1038/sj.onc.1207768>
65. Costello PS, Walters AE, Mee PJ, Turner M, Reynolds LF, Prisco A, Sarner N, Zamoyska R, Tybulewicz VL. The Rho-family GTP exchange factor Vav is a critical transducer of T cell receptor signals to the calcium, ERK, and NF-kappaB pathways. *Proc Natl Acad Sci USA* 1999; 96:3035-40; PMID:10077632; <http://dx.doi.org/10.1073/pnas.96.6.3035>
66. Reynolds LF, de Bertignies C, Norton T, Beeser A, Chernoff J, Tybulewicz VL. Vav1 transduces T cell receptor signals to the activation of the RasERK pathway via LAT, Sos, and RasGRP1. *J Biol Chem* 2004; 279:18239-46; PMID:14764585; <http://dx.doi.org/10.1074/jbc.M400257200>
67. Colucci F, Rosmaraki E, Bregenholt S, Samson SI, Di Bartolo V, Turner M, Vanes L, Tybulewicz V, Di Santo JP. Functional dichotomy in natural killer cell signaling: Vav1-dependent and -independent mechanisms. *J Exp Med* 2001; 193:1413-24; PMID:11413196; <http://dx.doi.org/10.1084/jem.193.12.1413>
68. Manetz TS, Gonzalez-Espinosa C, Arudchandran R, Xirasagar S, Tybulewicz V, Rivera J. Vav1 regulates phospholipase cgamma activation and calcium responses in mast cells. *Mol Cell Biol* 2001; 21:3763-74; PMID:11340169; <http://dx.doi.org/10.1128/MCB.21.11.3763-3774.2001>
69. Faccio R, Teitelbaum SL, Fujikawa K, Chappell J, Zal-lone A, Tybulewicz VL, Ross FP, Swat W. Vav3 regulates osteoclast function and bone mass. *Nat Med* 2005; 11:284-90; PMID:15711558; <http://dx.doi.org/10.1038/nm1194>
70. Sauzeau V, Sevilla MA, Montero MJ, Bustelo XR. The RhoRac exchange factor Vav2 controls nitric oxide-dependent responses in mouse vascular smooth muscle cells. *J Clin Invest* 2010; 120:315-30; PMID:20038798; <http://dx.doi.org/10.1172/JCI38356>
71. Norman KR, Fazio RT, Mellem JE, Espelt MV, Strange K, Beckerle MC, Mariq AV. The RhoRac-family guanine nucleotide exchange factor VAV-1 regulates rhythmic behaviors in *C. elegans*. *Cell* 2005; 123:119-32; PMID:16213217; <http://dx.doi.org/10.1016/j.cell.2005.08.001>
72. Moores SL, Selfors LM, Fredericks J, Breit T, Fujikawa K, Alt FW, Brugge JS, Swat W. Vav family proteins couple to diverse cell surface receptors. *Mol Cell Biol* 2000; 20:6364-73; PMID:10938113; <http://dx.doi.org/10.1128/MCB.20.17.6364-6373.2000>
73. Rodrigues L, Pires de Miranda M, Caloca MJ, Bustelo XR, Simas JP. Activation of Vav by the gammaherpesvirus M2 protein contributes to the establishment of viral latency in B lymphocytes. *J Virol* 2006; 80:6123-35; PMID:16731951; <http://dx.doi.org/10.1128/JVI.02700-05>
74. Gomez TS, Hamann MJ, McCarney S, Savoy DN, Lubking CM, Heldebrandt MP, Labno CM, McKean DJ, McNiven MA, Burkhardt JK, et al. Dynamin 2 regulates T cell activation by controlling actin polymerization at the immunological synapse. *Nat*

- Immunol 2005; 6:261-70; PMID:15696170; <http://dx.doi.org/10.1038/n1168>
75. Malartre M, Ayaz D, Amador FF, Martin-Bermudo MD. The guanine exchange factor vav controls axon growth and guidance during *Drosophila* development. *J Neurosci* 2010; 30:2257-67; PMID:20147552; <http://dx.doi.org/10.1523/JNEUROSCI.1820-09.2010>
76. Fernandez-Espartero CH, Ramel D, Farago M, Malartre M, Luque CM, Limanovich S, Katzav S, Emery G, Martin-Bermudo MD. GTP exchange factor Vav regulates guided cell migration by coupling guidance receptor signalling to local Rac activation. *J Cell Sci* 2013; 126:2285-93; PMID:23525006; <http://dx.doi.org/10.1242/jcs.124438>
77. Ruiz S, Santos E, Bustelo XR. The use of knockout mice reveals a synergistic role of the Vav1 and Rasgrf2 gene deficiencies in lymphomagenesis and metastasis. *PloS One* 2009; 4:e8229; PMID:20011522; <http://dx.doi.org/10.1371/journal.pone.0008229>
78. Sauzeau V, Jerkic M, Lopez-Novoa JM, Bustelo XR. Loss of Vav2 proto-oncogene causes tachycardia and cardiovascular disease in mice. *Mol Biol Cell* 2007; 18:943-52; PMID:17202406; <http://dx.doi.org/10.1091/mbc.E06-09-0877>
79. Quevedo C, Sauzeau V, Menacho-Marquez M, Castro-Castro A, Bustelo XR. Vav3-deficient mice exhibit a transient delay in cerebellar development. *Mol Biol Cell* 2011; 21:1125-39; PMID:20089829; <http://dx.doi.org/10.1091/mbc.E09-04-0292>
80. Sauzeau V, Horta-Junior JAC, Riobolobos AS, Fernandez G, Sevilla MA, Lopez DE, Montero MJ, Rico B, Bustelo XR. Vav3 is involved in GABAergic axon guidance events important for the proper function of brainstem neurons controlling cardiovascular, respiratory, and renal parameters. *Mol Biol Cell* 2010; 21:4251-63; PMID:20926682; <http://dx.doi.org/10.1091/mbc.E10-07-0639>
81. Guyenet PG. The sympathetic control of blood pressure. *Nat Rev Neurosci* 2006; 7:335-46; PMID:16760914; <http://dx.doi.org/10.1038/nrn1902>
82. Sauzeau V, Sevilla MA, Rivas-Elena JV, de Alava E, Montero MJ, Lopez-Novoa JM, Bustelo XR. Vav3 proto-oncogene deficiency leads to sympathetic hyperactivity and cardiovascular dysfunction. *Nat Med* 2006; 12:841-5; PMID:16767097; <http://dx.doi.org/10.1038/nm1426>
83. Menacho-Marquez M, Nogueiras R, Fabbiano S, Sauzeau V, Al-Massadi O, Dieguez C, Bustelo XR. Chronic sympathoexcitation through loss of Vav3, a Rac1 activator, results in divergent effects on metabolic syndrome and obesity depending on diet. *Cell Metab* 2013; 18:199-211; PMID:23931752; <http://dx.doi.org/10.1016/j.cmet.2013.07.001>
84. Cowan CW, Shao YR, Sahin M, Shamah SM, Lin MZ, Greer PL, Gao S, Griffith EC, Brugge JS, Greenberg ME. Vav family GEFs link activated Ephs to endocytosis and axon guidance. *Neuron* 2005; 46:205-17; PMID:15848800; <http://dx.doi.org/10.1016/j.neuron.2005.03.019>
85. Hale CF, Dietz KC, Varela JA, Wood CB, Zirlin BC, Leverich LS, Greene RW, Cowan CW. Essential role for vav Guanine nucleotide exchange factors in brain-derived neurotrophic factor-induced dendritic spine growth and synapse plasticity. *J Neurosci* 2011; 31:12426-36; PMID:21880903; <http://dx.doi.org/10.1523/JNEUROSCI.0685-11.2011>
86. Hunter SG, Zhuang G, Brantley-Sieders D, Swat W, Cowan CW, Chen J. Essential role of Vav family guanine nucleotide exchange factors in EphA receptor-mediated angiogenesis. *Mol Cell Biol* 2006; 26:4830-42; PMID:16782872; <http://dx.doi.org/10.1128/MCB.02215-05>
87. Liu JY, Seno H, Miletic AV, Mills JC, Swat W, Stapenbeck TS. Vav proteins are necessary for correct differentiation of mouse cecal and colonic enterocytes. *J Cell Sci* 2009; 122:324-34; PMID:19139088; <http://dx.doi.org/10.1242/jcs.033720>
88. Ruiz S, Santos E, Bustelo XR. RasGRF2, a guanine nucleotide exchange factor for Ras GTPases, participates in T-cell signaling responses. *Mol Cell Biol* 2007; 27:8127-42; PMID:17923690; <http://dx.doi.org/10.1128/MCB.00912-07>
89. Lawson CD, Donald S, Anderson KE, Patton DT, Welch HC. P-Rex1 and Vav1 cooperate in the regulation of formyl-methionyl-leucyl-phenylalanine-dependent neutrophil responses. *J Immunol* 2011; 186:1467-76; PMID:21178006; <http://dx.doi.org/10.4049/jimmunol.1002738>
90. Lopez-Lago M, Lee H, Cruz C, Movilla N, Bustelo XR. Tyrosine phosphorylation mediates both activation and downmodulation of the biological activity of Vav. *Mol Cell Biol* 2000; 20:1678-91; PMID:10669745; <http://dx.doi.org/10.1128/MCB.20.5.1678-1691.2000>
91. Katzav S, Cleveland JL, Heslop HE, Pulido D. Loss of the amino-terminal helix-loop-helix domain of the vav proto-oncogene activates its transforming potential. *Mol Cell Biol* 1991; 11:1912-20; PMID:2005887
92. Ming W, Li S, Billadeau DD, Quilliam LA, Dinanuer MC. The Rac effector p67phox regulates phagocyte NADPH oxidase by stimulating Vav1 guanine nucleotide exchange activity. *Mol Cell Biol* 2007; 27:312-23; PMID:17060455; <http://dx.doi.org/10.1128/MCB.00985-06>
93. Moarefi I, LaFevre-Bernt M, Sicheri F, Huse M, Lee CH, Kuriyan J, Miller WT. Activation of the Src-family tyrosine kinase Hck by SH3 domain displacement. *Nature* 1997; 385:650-3; PMID:9024665; <http://dx.doi.org/10.1038/385650a0>
94. Bustelo XR, Ledbetter JA, Barbacid M. Product of vav proto-oncogene defines a new class of tyrosine protein kinase substrates. *Nature* 1992; 356:68-71; PMID:1311423; <http://dx.doi.org/10.1038/356068a0>
95. Margolis B, Hu P, Katzav S, Li W, Oliver JM, Ullrich A, Weiss A, Schlessinger J. Tyrosine phosphorylation of vav proto-oncogene product containing SH2 domain and transcription factor motifs. *Nature* 1992; 356:71-4; PMID:1531699; <http://dx.doi.org/10.1038/356071a0>
96. Kant S, Swat W, Zhang S, Zhang ZY, Neel BG, Flavell RA, Davis RJ. TNF-stimulated MAP kinase activation mediated by a Rho family GTPase signaling pathway. *Genes Dev* 2011; 25:2069-78; PMID:21979919; <http://dx.doi.org/10.1101/gad.17224711>
97. Menacho-Marquez M, Garcia-Escudero R, Ojeda V, Abad A, Delgado P, Costa C, Ruiz S, Alarcon B, Paramio JM, Bustelo XR. The Rho exchange factors Vav2 and Vav3 favor skin tumor initiation and promotion by engaging extracellular signaling loops. *PLoS Biol* 2013; 11:e1001615; PMID:23935450; <http://dx.doi.org/10.1371/journal.pbio.1001615>
98. Gavard J, Gurtkind JS. VEGF controls endothelial-cell permeability by promoting the beta-arrestin-dependent endocytosis of VE-cadherin. *Nat Cell Biol* 2006; 8:1223-34; PMID:17060906; <http://dx.doi.org/10.1038/ncb1486>
99. Valls G, Codina M, Miller RK, Del Valle-Perez B, Vinyoles M, Caelles C, McCrea PD, Garcia de Herberos A, Dunach M. Upon Wnt stimulation, Rac1 activation requires Rac1 and Vav2 binding to p120-catenin. *J Cell Sci* 2012; 125:5288-301; PMID:22946057; <http://dx.doi.org/10.1242/jcs.101030>
100. Weng WK, Jarvis L, LeBien TW. Signaling through CD19 activates Vavmitogen-activated protein kinase pathway and induces formation of a CD19Vavphosphatidylinositol 3-kinase complex in human B cell precursors. *J Biol Chem* 1994; 269:32514-21; PMID:7528218
101. O'Rourke LM, Tooze R, Turner M, Sandoval DM, Carter RH, Tybulewicz VL, Fearon DT. CD19 as a membrane-anchored adaptor protein of B lymphocytes: costimulation of lipid and protein kinases by recruitment of Vav. *Immunity* 1998; 8:635-45; PMID:9620684; [http://dx.doi.org/10.1016/S1074-7613\(00\)80568-3](http://dx.doi.org/10.1016/S1074-7613(00)80568-3)
102. Son M, Park I, Lee OH, Rhee I, Park C, Yun Y. LIME mediates immunological synapse formation through activation of VAV. *Mol Cells* 2012; 33:407-14; PMID:22395814; <http://dx.doi.org/10.1007/s10059-012-0011-8>
103. Johmura S, Oh-hora M, Inabe K, Nishikawa Y, Hayaishi K, Vigorito E, Kitamura D, Turner M, Shingu K, Hikida M, et al. Regulation of Vav localization in membrane rafts by adaptor molecules Grb2 and BLNK. *Immunity* 2003; 18:777-87; PMID:12818159; [http://dx.doi.org/10.1016/S1074-7613\(03\)00139-0](http://dx.doi.org/10.1016/S1074-7613(03)00139-0)
104. Wu J, Motto DG, Koretzky GA, Weiss A. Vav and SLP-76 interact and functionally cooperate in IL-2 gene activation. *Immunity* 1996; 4:593-602; PMID:8673706; [http://dx.doi.org/10.1016/S1074-7613\(00\)80485-9](http://dx.doi.org/10.1016/S1074-7613(00)80485-9)
105. Tuosto L, Michel F, Acuto O. p95vav associates with tyrosine-phosphorylated SLP-76 in antigen-stimulated T cells. *J Exp Med* 1996; 184:1161-6; PMID:9064333; <http://dx.doi.org/10.1084/jem.184.3.1161>
106. Lindholm CK, Henriksson ML, Hallberg B, Welsh M. Shb links SLP-76 and Vav with the CD3 complex in Jurkat T cells. *Eur J Biochem* 2002; 269:3279-88; PMID:12084069; <http://dx.doi.org/10.1046/j.1432-1033.2002.03008.x>
107. Pauker MH, Hassan N, Noy E, Reicher B, Bardas-Saad M. Studying the dynamics of SLP-76, Nck, and Vav1 multimolecular complex formation in live human cells with triple-color FRET. *Sci Sig* 2012; 5:rs3; PMID:22534133; <http://dx.doi.org/10.1126/scisignal.2002423>
108. Bardas-Saad M, Shirasu N, Pauker MH, Hassan N, Perl O, Balbo A, Yamaguchi H, Houtman JC, Appella E, Schuck P, et al. Cooperative interactions at the SLP-76 complex are critical for actin polymerization. *EMBO J* 2010; 29:2315-28; PMID:20562827; <http://dx.doi.org/10.1038/emboj.2010.133>
109. Charvet C, Canonigo AJ, Billadeau DD, Altman A. Membrane localization and function of Vav3 in T cells depend on its association with the adapter SLP-76. *J Biol Chem* 2005; 280:15289-99; PMID:15708849; <http://dx.doi.org/10.1074/jbc.M500275200>
110. Zhang W, Irvin BJ, Tribble RP, Abraham RT, Samelson LE. Functional analysis of LAT in TCR-mediated signaling pathways using a LAT-deficient Jurkat cell line. *Int Immunol* 1999; 11:943-50; PMID:10360968; <http://dx.doi.org/10.1093/intimm/11.6.943>
111. Lahesmaa R, Allsup A, Soderberg C, Jackman J, Findell P, Peltz G. Modulation of the Grb2-associated protein complex in human CD4+ T cells by receptor activation. *J Immunol* 1995; 155:3815-22; PMID:7561087
112. Blancher F, Cardona A, Letimier FA, Hershsfeld MS, Acuto O. CD28 costimulatory signal induces protein arginine methylation in T cells. *J Exp Med* 2005; 202:371-7; PMID:16061726; <http://dx.doi.org/10.1084/jem.20050176>
113. Choudhary C, Kumar C, Gnani F, Nielsen ML, Rehman M, Walther TC, Olsen JV, Mann M. Lysine acetylation targets protein complexes and co-regulates major cellular functions. *Science* 2009; 325:834-40; PMID:19608861; <http://dx.doi.org/10.1126/science.1175371>
114. Puga I, Rao A, Macian F. Targeted cleavage of signaling proteins by caspase 3 inhibits T cell receptor signaling in anergic T cells. *Immunity* 2008; 29:193-204; PMID:18701083; <http://dx.doi.org/10.1016/j.immuni.2008.06.010>
115. Hofmann TG, Hehner SP, Droge W, Schmitz ML. Caspase-dependent cleavage and inactivation of the Vav1 proto-oncogene product during apoptosis prevents IL-2 transcription. *Oncogene* 2000; 19:1153-63; PMID:10713703; <http://dx.doi.org/10.1038/sj.onc.1203406>
116. Razidlo GL, Wang Y, Chen J, Krueger EW, Billadeau DD, McNiven MA. Dynamin 2 potentiates invasive migration of pancreatic tumor cells through



- stabilization of the Rac1 GEF Vav1. *Dev Cell* 2013; 24:573-85; PMID:23537630; <http://dx.doi.org/10.1016/j.devcel.2013.02.010>
117. Yoo AS, Greenwald I. LIN-12/Notch activation leads to microRNA-mediated down-regulation of Vav in *C. elegans*. *Science* 2005; 310:1330-3; PMID:16239437; <http://dx.doi.org/10.1126/science.1119481>
118. Wang R, Zhao N, Li S, Fang JH, Chen MX, Yang J, Jia WH, Yuan Y, Zhuang SM. MicroRNA-195 suppresses angiogenesis and metastasis of hepatocellular carcinoma by inhibiting the expression of VEGF, VAV2, and CDC42. *Hepatology* 2013; 58:642-53; PMID:23468064; <http://dx.doi.org/10.1002/hep.26373>
119. Huang X, Ding L, Bennetwih KL, Tong RT, Welford SM, Ang KK, Story M, Le QT, Giaccia AJ. Hypoxia-inducible mir-210 regulates normoxic gene expression involved in tumor initiation. *Mol Cell* 2009; 35:856-67; PMID:19782034; <http://dx.doi.org/10.1016/j.molcel.2009.09.006>
120. Fernandez-Zapico ME, Gonzalez-Paz NC, Weiss E, Savoy DN, Molina JR, Fonseca R, Smyrk TC, Chari ST, Urrutia R, Billadeau DD. Ectopic expression of VAV1 reveals an unexpected role in pancreatic cancer tumorigenesis. *Cancer Cell* 2005; 7:39-49; PMID:15652748; <http://dx.doi.org/10.1016/j.ccr.2004.11.024>
121. Carvajal-Gonzalez JM, Mulero-Navarro S, Roman AC, Sauzeau V, Merino JM, Bustelo XR, Fernandez-Salguero PM. The dioxin receptor regulates the constitutive expression of the vav3 proto-oncogene and modulates cell shape and adhesion. *Mol Biol Cell* 2009; 20:1715-27; PMID:19158396; <http://dx.doi.org/10.1091/mbc.E08-05-0451>
122. Ilan L, Katzav S. Human Vav1 expression in hematopoietic and cancer cell lines is regulated by c-Myb and by CpG methylation. *PLoS One* 2012; 7:e29939; PMID:22253833; <http://dx.doi.org/10.1371/journal.pone.0029939>
123. Sauzeau V, Carvajal-Gonzalez JM, Rioloobos AS, Sevilla MA, Menacho-Marquez M, Roman AC, Abad A, Montero MJ, Fernandez-Salguero P, Bustelo XR. Transcriptional factor aryl hydrocarbon receptor (Ahr) controls cardiovascular and respiratory functions by regulating the expression of the Vav3 proto-oncogene. *J Biol Chem* 2011; 286:2896-909; PMID:21115475; <http://dx.doi.org/10.1074/jbc.M110.187534>
124. Gulbins E, Coggeshall KM, Baier G, Telford D, Langlet C, Baier-Bitterlich G, Bonnefoy-Berard N, Burn P, Wittinghofer A, Altman A. Direct stimulation of Vav guanine nucleotide exchange activity for Ras by phorbol esters and diglycerides. *Mol Cell Biol* 1994; 14:4749-58; PMID:7516472
125. Han J, Luby-Phelps K, Das B, Shu X, Xia Y, Mosteller RD, Krishna UM, Falck JR, White MA, Broek D. Role of substrates and products of PI 3-kinase in regulating activation of Rac-related guanine triphosphatases by Vav. *Science* 1998; 279:558-60; PMID:9438848; <http://dx.doi.org/10.1126/science.279.5350.558>
126. Kazanietz MG, Bustelo XR, Barbacid M, Kolch W, Mischak H, Wong G, Pettit GR, Bruns JD, Blumberg PM. Zinc finger domains and phorbol ester pharmacophore. Analysis of binding to mutated form of protein kinase C zeta and the vav and c-raf proto-oncogene products. *J Biol Chem* 1994; 269:11590-4; PMID:8157692
127. Geczy T, Peach ML, El Kazzouli S, Sigano DM, Kang JH, Valle CJ, Selezneva J, Woo W, Keddi N, Lewin NE, et al. Molecular basis for failure of "atypical" C1 domain of Vav1 to bind diacylglycerolphorbol ester. *J Biol Chem* 2012; 287:13137-58; PMID:22351766; <http://dx.doi.org/10.1074/jbc.M111.320010>
128. Park WS, Heo WD, Whalen JH, O'Rourke NA, Bryan HM, Meyer T, Teruel MN. Comprehensive identification of PIP3-regulated PH domains from *C. elegans* to *H. sapiens* by model prediction and live imaging. *Mol Cell* 2008; 30:381-92; PMID:18471983; <http://dx.doi.org/10.1016/j.molcel.2008.04.008>
129. Isakoff SJ, Cardozo T, Andreev J, Li Z, Ferguson KM, Abagyan R, Lemmon MA, Aronheim A, Skolnik EY. Identification and analysis of PH domain-containing targets of phosphatidylinositol 3-kinase using a novel in vivo assay in yeast. *EMBO J* 1998; 17:5374-87; PMID:9736615; <http://dx.doi.org/10.1093/emboj/17.18.5374>
130. Inabe K, Ishiai M, Scharenberg AM, Freshney N, Downward J, Kurosaki T. Vav3 modulates B cell receptor responses by regulating phosphoinositide 3-kinase activation. *J Exp Med* 2002; 195:189-200; PMID:11805146; <http://dx.doi.org/10.1084/jem.20011571>
131. Bassermann F, Jahn T, Miething C, Seipel P, Bai RY, Coutinho S, Tybulewicz VL, Peschel C, Duyster J. Association of Bcr-Abl with the proto-oncogene Vav is implicated in activation of the Rac-1 pathway. *J Biol Chem* 2002; 277:12437-45; PMID:11790798; <http://dx.doi.org/10.1074/jbc.M112397200>
132. Harnois T, Constantin B, Rioux A, Grenioux E, Kitzis A, Bourmeyster N. Differential interaction and activation of Rho family GTPases by p210bcr-abl and p190bcr-abl. *Oncogene* 2003; 22:6445-54; PMID:14508524; <http://dx.doi.org/10.1038/sj.onc.1206626>
133. Patel V, Rosenfeldt HM, Lyons R, Servitja JM, Bustelo XR, Siroff M, Gutkind JS. Persistent activation of Rac1 in squamous carcinomas of the head and neck: evidence for an EGFRVav2 signaling axis involved in cell invasion. *Carcinogenesis* 2007; 28:1145-52; PMID:17234718; <http://dx.doi.org/10.1093/carcin/bgm008>
134. Levaot N, Voytyuk O, Dimitriou I, Sircoulomb F, Chandrakumar A, Deckert M, Krzyzanowski PM, Scott A, Gu S, Janmohamed S, et al. Loss of Tankyrase-mediated destruction of 3BP2 is the underlying pathogenic mechanism of cherubism. *Cell* 2011; 147:1324-39; PMID:22153076; <http://dx.doi.org/10.1016/j.cell.2011.10.045>
135. Pires de Miranda M, Alenquer M, Marques S, Rodrigues L, Lopes F, Bustelo XR, Simas JP. The Gammaherpesvirus m2 Protein Manipulates the Fyn-Vav Pathway through a Multidocking Mechanism of Assembly. *PLoS One* 2008; 3:e1654; PMID:18301737; <http://dx.doi.org/10.1371/journal.pone.0001654>
136. Pires de Miranda M, Lopes FB, McVey CE, Bustelo XR, Simas JP. Role of Src homology domain binding in signaling complexes assembled by the murine gamma-herpesvirus M2 protein. *J Biol Chem* 2013; 288:3858-70; PMID:23258536; <http://dx.doi.org/10.1074/jbc.M112.439810>
137. Jagodic M, Colacios C, Nohra R, Dejean AS, Beyeen AD, Khademi M, Casemayou A, Lamouroux L, Duthoit C, Papapietro O, et al. A role for VAV1 in experimental autoimmune encephalomyelitis and multiple sclerosis. *Sci Transl Med* 2009; 1:10ra21; PMID:20368159; <http://dx.doi.org/10.1126/scitranslmed.3000278>
138. Aleksic B, Kushima I, Hashimoto R, Ohi K, Ikeda M, Yoshimi A, Nakamura Y, Ito Y, Okochi T, Fukuo Y, et al. Analysis of the VAV3 as candidate gene for schizophrenia: evidences from voxel-based morphometry and mutation screening. *Schizophr Bull* 2013; 39:720-8; PMID:22416266; <http://dx.doi.org/10.1093/schbul/sbs038>
139. Eriksson N, Tung JY, Kiefer AK, Hinds DA, Francke U, Mountain JL, Do CB. Novel associations for hypothyroidism include known autoimmune risk loci. *PLoS One* 2012; 7:e34442; PMID:22493691; <http://dx.doi.org/10.1371/journal.pone.0034442>
140. Fujikawa K, Iwata T, Inoue K, Akahori M, Kadotani H, Fukaya M, Watanabe M, Chang Q, Barnett EM, Swat W. VAV2 and VAV3 as candidate disease genes for spontaneous glaucoma in mice and humans. *PLoS One* 2010; 5:e9050; PMID:20140222; <http://dx.doi.org/10.1371/journal.pone.0009050>
141. Shi D, Takano Y, Nakazawa T, Mengkegale M, Yokokura S, Nishida K, Fuse N. Molecular genetic analysis of primary open-angle glaucoma, normal tension glaucoma, and developmental glaucoma for the VAV2 and VAV3 gene variants in Japanese subjects. *Biochem Biophys Res Comm* 2013; 432:509-12; PMID:23402756; <http://dx.doi.org/10.1016/j.bbrc.2013.02.003>
142. Rao KN, Kaur I, Parikh RS, Mandal AK, Chandrasekhar G, Thomas R, Chakrabarti S. Variations in NTF4, VAV2, and VAV3 genes are not involved with primary open-angle and primary angle-closure glaucomas in an Indian population. *Invest Ophthalmol Vis Sci* 2010; 51:4937-41; PMID:20463313; <http://dx.doi.org/10.1167/iovs.10-5553>
143. Martin H, Mali RS, Ma P, Chatterjee A, Ramdas B, Sims E, Munugalavada V, Ghosh J, Mattingly RR, Visconte V, et al. Pak and Rac GTPases promote oncogenic KIT-induced neoplasms. *J Clin Invest* 2013; 123:4449-63; PMID:24091327; <http://dx.doi.org/10.1172/JCI67509>
144. Chang KH, Sanchez-Aguilera A, Shen S, Sengupta A, Madhu MN, Ficker AM, Dunn SK, Kuenzi AM, Arnett JL, Santho RA, et al. Vav3 collaborates with p190-Bcr-ABL in lymphoid progenitor leukemogenesis, proliferation and survival. *Blood* 2012; 120:800-11; PMID:22692505; <http://dx.doi.org/10.1182/blood-2011-06-361709>
145. Rahaman SO, Li W, Silverstein RL. Vav Guanine nucleotide exchange factors regulate atherosclerotic lesion development in mice. *Arterioscler Thromb Vasc Biol* 2013; 33:2053-7; PMID:23825362; <http://dx.doi.org/10.1161/ATVBAHA.113.301414>
146. Chen K, Li W, Major J, Rahaman SO, Febbraio M, Silverstein RL. Vav guanine nucleotide exchange factors link hyperlipidemia and a prothrombotic state. *Blood* 2011; 117:5744-50; PMID:21427288; <http://dx.doi.org/10.1182/blood-2009-01-201970>
147. Rahaman SO, Swat W, Febbraio M, Silverstein RL. Vav family Rho guanine nucleotide exchange factors regulate CD36-mediated macrophage foam cell formation. *J Biol Chem* 2011; 286:7010-7; PMID:21209086; <http://dx.doi.org/10.1074/jbc.M110.192450>
148. Haubert D, Li J, Saveliev A, Calzascia T, Sutter E, Metzler B, Kaiser D, Tybulewicz VL, Weckbecker G. Vav1 GEF activity is required for T cell mediated allograft rejection. *Transpl Immunol* 2012; 26:212-9; PMID:22456277; <http://dx.doi.org/10.1016/j.trim.2012.03.003>
149. Yi F, Xia M, Li N, Zhang C, Tang L, Li PL. Contribution of guanine nucleotide exchange factor Vav2 to hyperhomocysteinemic glomerulosclerosis in rats. *Hypertension* 2009; 53:90-6; PMID:19029489; <http://dx.doi.org/10.1161/HYPERTENSIONAHA.108.115675>
150. Andre G, Sandoval JE, Retailleau K, Loufrani L, Toumaniantz G, Offermanns S, Rolli-Derkinderen M, Loirand G, Sauzeau V. Smooth muscle specific Rac1 deficiency induces hypertension by preventing p116RIP3-dependent RhoA inhibition. *J Am Heart Ass* 2014; 3:e000852; PMID:24938713; <http://dx.doi.org/10.1161/JAHA.114.000852>
151. Fabbiano S, Menacho-Márquez M, Sevilla MA, Albarrán-Juárez J, Zheng Y, Offermanns S, Montero MJ, Bustelo XR. Genetic dissection of the Vav2-Rac1 signaling axis in vascular smooth muscle cells. *Mol Cell Biol* 2014; PMID:25288640; <http://dx.doi.org/10.1128/MCB.01066-14>
152. Vigil D, Cherfils J, Rossman KL, Der CJ. Ras superfamily GEFs and GAPs: validated and tractable targets for cancer therapy? *Nat Rev Cancer* 2010; 10:842-57; PMID:21102635; <http://dx.doi.org/10.1038/nrc2960>
153. Shang X, Marchionni F, Evelyn CR, Sipes N, Zhou X, Seibel W, Wortman M, Zheng Y. Small-molecule inhibitors targeting G-protein-coupled Rho guanine nucleotide exchange factors. *Proc Natl Acad Sci USA* 2013; 110:3155-60; PMID:23382194; <http://dx.doi.org/10.1073/pnas.1212341110>